

phenylmethylsulfonyl fluoride/ protease inhibitor cocktail (Sigma)), and centrifuged at 12000 x g at 4°C for 15 minutes. Supernatants containing soluble proteins were stored at -80°C. Protein concentrations were determined using a BCA kit (Pierce, Rockford, IL). Western blotting analysis was performed as described (Lan et al., 1999) using either SDS 15% PAGE or SDS 4-20% gradient PAGE (ISC BioExpress, Kaysville, UT). The protein blots were incubated with 1:1500 dilution of rabbit polyclonal anti-AeSCP-2 antibodies. The goat anti-rabbit horseradish peroxidase conjugated secondary antibody (Jackson ImmunoResearch laboratory, West Grove, PA) was used at 1:3300 dilution. DAB solution (0.3 mg/ml and 0.03% hydrogen peroxide in PBS) was used to visualize the bound antibodies, which was developed within 5 minutes at room temperature.

In the specification at page 44, please replace line ~~21~~²² with the following line:

BC
9/19/06

FlyBase. 2001. [[http://]]flybase.bio.indiana.edu:82[[/]]

In the specification at page 47, please replace line 14 with the following line:

NCBI, 2002. [[http://]]www.ncbi.nlm.nih.gov/Structure/lexington/lexington.cgi